OM of: US-08-962-560A-4 to: N_Geneseq_36:*   out_format : pfs

Date: Feb 5, 2000   8:38 AM

About: Results were produced by the GenCore software, version 4.5,
          Copyright (c) 1993-2000 Compugen Ltd.

Command line parameters:
-MODEL=frame+_p2n.model -DEV-xlp
-Q=/cgnl_1/USPTO_spool/US08962560/runat_03022000_141841_3613/app_query.fasta.1
-DB-N_Geneseq_36 -QFMT=fastap -SUFFIX=rng -GAPOP=12.000
-GAPEXT=4.000 -MINMATCH-0.100 -LOOPCL=0.000 -LOOPEXN=0.000
-QGAPOP-4.500 -QGAPEXN=0.050 -XGAPEXN-0.500
-FGAPOP-6.000 -FGAPEXT-7.000 -YGAPOP=10.000 -YGAPEXN-0.500
-DELOP-6.000 -DELEXT=7.000 -START=1 -MATRIX=blosum62
-TRANS-human40.cdi -LIST-45 -DOCALIGN=200 -THR_SCORE=pct
-ALIGN-15 -MODE=LOCAL -OUTFMT=pfs -NORM=ext -MINLEN=0
-MAXLEN-1000000 -USER=US08962560 -NCPU-6 -ICPU-3 -NO_XLPXY -WAIT
-THREADS-1

Search Information block:
Query: US-08-962-560A-4
Query length: 212
Database: N_Geneseq_36:*
Database sequences: 311585
Database length: 125090042
Search time (sec): 1063.640000

score_list:

Sequence	Strd	Orig	ZScore	EScore	Len	! Documentation
N_Geneseq_36:V38659	+	1107.00	1425.53	8.4e-72	1235	! Mus musculus SOCS1 gene. Supp
N_Geneseq_36:V42701	+	1093.00	1408.54	7.4e-71	1087	! cDNA encoding a STAT function
N_Geneseq_36:V38663	+	1073.00	1374.61	5.8e-69	2807	! Rattus norvegicus SOCS1 gene.
N_Geneseq_36:V38662	+	1053.50	1357.47	5.2e-68	1094	! Homo sapiens SOCS1 gene. Supp
N_Geneseq_36:T43380	+	249.50	314.06	6.8e-10	1960	! Human cytokine response gene C
N_Geneseq_36:V38660	+	234.50	299.46	4.4e-09	1121	! Mus musculus SOCS2 gene. Suppr
N_Geneseq_36:V69307	+	234.50	293.17	9.9e-09	2342	! Human EPRG1 cDNA #1. New EPO F
N_Geneseq_36:V69309	+	234.50	293.17	9.9e-09	2342	! Human EPRG1 cDNA derived from
N_Geneseq_36:T96002	+	297.07	297.07	6.0e-09	1374	! Human cytokine-inducible SH2-c
N_Geneseq_36:V38861	+	220.50	287.29	2.1e-08	2187	! Mus musculus SOCS3 gene. Suppr
N_Geneseq_36:V34148	+	221.00	275.60	9.5e-08	2378	! Human secreted protein gene 35
N_Geneseq_36:V38681	+	147.50	189.47	0.0059	848	! Homo sapiens SOCS11 gene. Supp
N_Geneseq_36:V38686	+	145.00	183.13	0.0134	1221	! Homo sapiens SOCS5 gene. Suppr
N_Geneseq_36:V38687	+	138.50	168.83	0.0838	2438	! Mus musculus SOCS14 cDNA. Supp
N_Geneseq_36:V71075	+	126.50	153.01	0.6374	2532	! Green fluorescent protein-prot
N_Geneseq_36:V38867	+	126.00	152.35	0.6939	2537	! Mus musculus SOCS5 gene. Suppr
N_Geneseq_36:Q21065	+	129.50	129.64	12.77	17041	! Genomic DNA of human PACAP 38
N_Geneseq_36:Q67057	+	121.00	129.64	12.77	17041	! PACAP38 DNA. PACAP38 gene pr
N_Geneseq_36:V71026	+	119.50	142.79	2.36	2907	! p85alpha-green flourescent pro
N_Geneseq_36:V71035	+	119.50	142.77	2.37	2913	! p85alpha-green flourescent pro
N_Geneseq_36:Q27254	+	119.50	141.53	2.78	3372	! Growth Factor Receptor Bound F
N_Geneseq_36:T01198	+	119.50	141.53	2.78	3372	! Human GRB-1 genomic DNA sequen
N_Geneseq_36:T35508	+	118.50	145.85	1.60	1747	! Rat M1PTP gene cDNA. clone der
N_Geneseq_36:T12235	+	118.50	135.70	5.87	5737	! pp60PIK cDNA. Pp60PIK peptide
N_Geneseq_36:V38677	+	116.50	141.37	2.84	5737	! Homo sapiens SOCS9 gene. Suppr
N_Geneseq_36:T80413	-	116.50	115.87	74.67	43280	! Tylactone synthase gene clust
N_Geneseq_36:X39794	-	149.00	149.68	0.9778	764	! Gastric cancer associated gene.
N_Geneseq_36:V71026	-	113.50	136.12	5.56	2562	! Protein tyrosine phosphatase 1
N_Geneseq_36:Q48806	-	109.50	109.99	158.71	29879	! eryA region of S. erythraea c
N_Geneseq_36:T43076	-	109.00	128.10	15.56	3319	! Wheat acetyl-CoA carboxylase A
N_Geneseq_36:V33412	+	109.00	128.10	15.56	3319	! 5' end flanking sequence of wh
N_Geneseq_36:V09214	+	107.00	130.36	11.64	1881	! CrkI cDNA sequence. Use of ant
N_Geneseq_36:Q06844	+	107.00	128.75	14.31	2271	! Amylase gene from Streptomyces
N_Geneseq_36:Q29515	-	107.00	125.36	22.12	3382	! Bovine herpes virus type 1 gly
N_Geneseq_36:Q48849	-	107.00	125.36	22.12	3382	! Bovine herpesvirus type 1 gi g
N_Geneseq_36:Q67058	-	105.50	137.40	4.72	657	! PACPAP38 promoter. PACAP38 gene
N_Geneseq_36:V09005	-	105.50	114.99	83.57	9077	! Vector containing human TGFbe
N_Geneseq_36:V09006	-	105.50	114.99	83.57	9077	! Vector containing Rabbit TGFbe
N_Geneseq_36:T35310	-	104.50	128.39	15.00	1624	! Human SH-PTP1 cDNA clone deriv
N_Geneseq_36:T35306	-	104.50	126.01	20.34	2145	! Human SH-PTP1 cDNA (protein ty
N_Geneseq_36:N80309	+	103.00	123.51	28.02	2291	! Entire amylase gene. Expressid

N_Geneseq_36:T97129	+	102.50	124.70	24.06	1848	! Human netrin-1 cDNA. Human
N_Geneseq_36:V21187	+	102.50	95.93	963.02	53789	! Amycolatopsis mediterranei
N_Geneseq_36:V34002	+	102.00	126.42	19.30	1401	! S. peucetius dnrX gene. Pro
N_Geneseq_36:V69308	+	101.00	121.63	35.68	2111	! Human EPRG1 cDNA #2. New EP

seq_name: N_Geneseq_36:V38659

seq_documentation_block:
ID   V38659 standard; DNA; 1235 BP.
AC   V38659;
DT   27-OCT-1998 (first entry)
DE   Mus musculus SOCS1 gene.
DE   Mus musculus SOCS1 gene.
KW   SOCS; suppressor of cytokine signalling; PCR primer;
KW   autoimmune disease; diagnosis; cancer; treatment;
KW   cytokine mediated cellular responsiveness; hyperimmunity;
KW   immunosuppression; allergies; hypertension; ss.
OS   Mus musculus.
OS   Mus musculus.
FH   Key   Location/Qualifiers
FT   CDS   161..799
FT   /*tag= a
FT   /product= SOCS1 protein
PN   WO9820023-A1.
PD   14-MAY-1998.
PF   31-OCT-1997; AU0729.
PF   14-FEB-1997; AU-005117.
PF   01-NOV-1996; AU-003384.
PS   Claim 14; Page 108-109; 32pp; English.
PT   The sequence is that of a gene encoding a suppressor of cytokine
PT   signalling protein (SOCS). SOCs can be used to screen for naturally
PT   occurring antibodies to SOCS, which may occur, e.g. in some autoimmune
PT   diseases. Alternatively, specific antibodies can be used to
PT   screen for SOCS, which is useful as a knowledge of SOCS levels
PT   may be important for the diagnosis of certain cancers. Soluble
PT   SOCS polypeptides can be used to treat disease, injury or
PT   abnormality involving cytokine mediated cellular responsiveness,
PT   e.g. hyperimmunity, immunosuppression, allergies and hypertension.
PI   Alexander WS, Hilton DJ, Metcalf D, Nicholson SE;
PI   Nicola NA, Richardson RT, Starr R, Viney EM, Willson TA;
PA   (HALL-) HALL INST MEDICAL RES WALTER & ELIZA.
PR   01-NOV-1996; AU-003384.
DR   WPI; 98-286884/25.
DR   P-PSDB; W62613.
SQ   Sequence 1235 BP; 421 C; 347 G; 192 A; 275 T;

seq_documentation_block:
seq_documentation_block:

Search Information block:

Suppressor of cytokine signalling proteins - useful to treat
disease, injury or abnormality involving cytokine mediated cellular
responsiveness e.g. hyperimmunity, immunosuppression, allergies and
hypertension

alignment_block:
US-08-962-560A-4 x V38659

alignment_scores:
          Quality: 1107.00   Length: 212
            Ratio: 5.222   Gaps: 0
   Percent Similarity: 100.000   Percent Identity: 100.000

Align seg 1/1   to: V38659   from: 1   to: 1235

```
    1 MetValAlaArgAsnGlnValAlaIleAlaAspAsnAlaIleSerProAlaAl 17
      ||||||||||||||||||||||||||||||||||||||||||||||||||||||
  161 ATGGTAGCACGCAACCAGGTGGCAGCCGACAATGCGATCTCCCGGCAGC 210

   17 aGluProArgArgAsrgSerGluProSerSerSerSerSerSerP 34
      ||||||||||||||||||||||||||||||||||||||||||||||
  211 AGAGCCCCGACGGCGGGTCAGAGCCCTCCTCGTCTTCGTCCTCGC 260

   34 roAlaAlaProValArgProArgProCysProLaVaProAlaProAla 50
      ||||||||||||||||||||||||||||||||||||||||||||||||
  261 CACCGGCCCCGTGGCGTCCCGCCCTGCCCCGGCGTCCCAGCCCCACC 310

   51 ProGlyAspThrHisPheArgThrPheArgSerHisSerAspTyrArgAr 67
      |||||||||||||||||||||||||||||||||||||||||||||||||
  311 CCTGGCGACACTCACTTCCGCACCTTCCGCTCCCACTCCGATACCGGCG 360
```

Left column

```
 67 glleThrArgThrSerAlaLeuLeuAspAlaCysGlyPheTyrTrpGlyP  84
    ||||||||||||||||||||||||||||||||||||||||||||||||||
361 CATCACCCGGACCAGCGCCCTCCTGGACGCTTGCGGGTTCTATTGGGGAC 410

 84 roLeuSerValHisGlyAlaHisGluArgLeuArgAlaGluProValGly 100
    ||||||||||||||||||||||||||||||||||||||||||||||||||
411 CCCTGAGCGTGCACGGGGCGCACGAGCGGCTGCGTGCCGAGCCGGTGGGC 460

101 ThrPheLeuValArgAspSerArgGlnArgAsnCysPhePheAlaLeuSe 117
    ||||||||||||||||||||||||||||||||||||||||||||||||||
461 ACCTTCTTGGTGCGCGACAGTCGTCAACGAGAACTGCTTCTTCGCGCTCAG 510

117 rValLysMetAlaSerGlyProThrSerIleArgValHisPheGlnAlaG 134
    ||||||||||||||||||||||||||||||||||||||||||||||||||
511 CGTGAAGATGGCTTCGGGCCCCACGAGCCATCCGGCGTGCACTTCCAGGCCG 560

134 lyArgPheHisLeuAspGlySerArgGluThrPheAspCysLeuPheGlu 150
    ||||||||||||||||||||||||||||||||||||||||||||||||||
561 GCCGGCTTCCACTTGGACGGCAGCCGGGAGACCTTCGACTGCCTTTTCGAG 610

151 LeuLeuGluHisTyrValAlaLeuAlaProArgMetLeuGlyAlaProIle 167
    ||||||||||||||||||||||||||||||||||||||||||||||||||
611 CTGCTGGAGCACTACGTGGCCCTGGCGCCCCGGATGTTGGGGGCCCCGATC 660

167 uArgGlnArgArgValArgProLeuGlnGluLeuCysSerArgGlnArgIleV 184
    ||||||||||||||||||||||||||||||||||||||||||||||||||
661 GCGCCAGCGCCGTCGGGCCCCTGCAGGAGCTGTGTCGCCAGGCGATCG 710

184 alAlaAlaValGlyArgGluAsnLeuAlaArgIleProLeuAsnProVal 200
    ||||||||||||||||||||||||||||||||||||||||||||||||||
711 TGGCCGCGGTCGGGCGAGAACCTGGCGCGCATCCCTCTTAACCCGGTA 760

201 LeuArgAspThrTyrLeuSerSerPhePheProGlnIle 212
    |||||||||||||||||||||||||||||||||||||
761 CTCCGTGACTACCTGAGTTCCTTCCCTTCCCAGATC 796
```

seq_name: N_Geneseq_36:V42701

seq_documentation_block:

```
ID   V42701 standard; cDNA; 1087 BP.
AC   V42701;
DT   30-OCT-1998 (first entry)
DE   cDNA encoding a STAT function regulatory protein designated SIIS-1.
KW   SIIS-1; STAT-induced inhibitor; STAT function;
KW   JAK/STAT signal transmission system; STAT3; STAT6; inhibit;
KW   tyrosine phosphorylation; gp130; cytokine-regulating protein; CIS;
KW   screen; cytokine regulatory; inhibitory activity; ds.
OS   Mus sp.
FH   Key             Location/Qualifiers
FT   CDS             16..654
FT                   /*tag= a
PI   Naka T;
PA   (KISH/) KISHIMOTO T.
PR   10-JAN-1997; JP-014737.
PF   23-OCT-1997; J03860.
PD   16-JUL-1998.
PN   WO9830688-A1.
DR   P-PSDB; W70962.
WPI  98-399137/34.
PT   STAT function regulatory protein - used in screening candidate
PT   substances for cytokine regulatory activity
PS   Claim 5; Pages 39-41; 60bp; Japanese.
CC   The present sequence encodes a protein (designated SIIS-1, STAT-induced
CC   inhibitor of STAT function 1) which regulates STAT protein function in
CC   the JAK/STAT signal transmission system in mammalian cells. The protein
CC   is induced by STAT3 or STAT6. It inhibits tyrosine phosphorylation of
CC   STAT3 and of gp130. The SIIS-1 protein sequence contains an SH2 domain
CC   and is related to the cytokine-regulating protein CIS. SIIS-1, or
CC   transformant cells expressing it, may be used to screen candidate
CC   substances for cytokine regulatory or inhibitory activity.
SQ   Sequence 1087 BP; 171 A; 363 C; 305 G; 248 T;
```

Right column

```
alignment_scores:
                Quality:  1093.00        Length:   212
                Ratio:       5.180       Gaps:       0
Percent Similarity:  99.528     Percent Identity:  98.585

alignment_block:
US-08-962-560A-4 x V42701

Align seg 1/1  to: V42701  from: 1  to: 1087
```

```
  1 MetValAlaArgAsnGlnValAlaAlaAspAsnAlaIleSerProAlaAl  17
    |||||||||||||||||||||||||||||||||||||||||||||||||||
  1 ATGGTAGCACGCAACCAGGTGGCAGCCGACAATGCGATCTCCCCGGCGAC  65

 17 aGluProArgArgArgSerGluSerProSerSerSerSerSerSerSerP  34
    ||||||||||||||||||||||||||||||||||||||||||||||||||
 66 AGAGCCCCGACGGGCGGGTCAGAGCCCTCCTCGTCCTCGTCCTCGTCGCGC 115

 34 roAlaAlaLaProValArgProArgProCysProAlaValProAlaProAla 50
    ||||||||||||||||||||||||||||||||||||||||||||||||||
116 CACGGCGCCCCTGCGTCCCCGGGGTCCCAGCCCCCAGCC 165

 51 ProGlyAspThrHisPheAspArgThrArgHisSerAspTyrTyrArgAr  67
    ||||||||||||||||||||||||||||||||||||||||||||||||||
166 CCTGGCGACACTCACTTCGGACCTTCCGGCTCCCCACTCCGATTACCGGCG 215

 67 glleThrArgThrSerAlaLeuLeuAspAlaCysGlyPheTyrTrpGlyP  84
    ||||||||||||||||||||||||||||||||||||||||||||||||||
216 CATCACCCGGACCAGCGCCCTCCTGGACGCTTGCGGGTTCTATTGGGGAC 265

 84 roLeuSerValHisGlyAlaHisGluArgLeuArgAlaGluProValGly 100
    ||||||||||||||||||||||||||||||||||||||||||||||||||
266 CCCTGAGCGTGCACGGGGCGCACGAGCGGCTGCGTGCCGAGCCGGTGGGC 315

101 ThrPheLeuValArgAspSerArgGlnArgAsnCysPhePheAlaLeuSe 117
    ||||||||||||||||||||||||||||||||||||||||||||||||||
316 ACCTTCTTGGTGCGCGACAGTCGTCAACGAGAACTGCTTCTTCGCGCTCAG 365

117 rValLysMetAlaSerGlyProThrSerIleArgValHisPheGlnAlaG 134
    ||||||||||||||||||||||||||||||||||||||||||||||||||
366 CGTGAAGATGGCTTCGGGCCCCACGAGCCATCCGGCGTGCACTTCCAGGCCG 415

134 lyArgPheHisLeuAspGlySerArgGluThrPheAspCysLeuPheGlu 150
    ||||||||||||||||||||||||||||||||||||||||||||||||||
416 GCCGGCTTCCACTTGGACGGCAGCCGGGAGACCTTCGACTGCCTTTTCGAG 465

151 LeuLeuGluHisTyrValAlaLeuAlaProArgMetLeuGlyAlaProIle 167
    ||||||||||||||||||||||||||||||||||||||||||||||||||
466 CTGCTGGAGCACTACGTGGCCCTGGCGCCCCGGATGTTGGGGGCCCCGATC 515

167 uArgGlnArgArgValArgProLeuGlnGluLeuCysSerArgGlnArgIleV 184
    ||||||||||||||||||||||||||||||||||||||||||||||||||
516 GCGCCAGCGCCGTCGGGCCCCTGCAGGAGCTGTGTCGCCAGGCGATCG 565

184 alAlaAlaValGlyArgGluAsnLeuAlaArgIleProLeuAsnProVal 200
    ||||||||||||||||||||||||||||||||||||||||||||||||||
566 TGGCCGCGGTCGGGCGAGAACCTGGCGCGCATCCCTCTTAACCCGGTA 615

201 LeuArgAspThrTyrLeuSerSerPhePheProGlnIle 212
    |||||||||||||||||||||||||||||||||||||
616 CTCCGTGACTACCTGAGTTCCTTCCCTTCCCAGATC 651
```

seq_name: N_Geneseq_36:V38663

seq_documentation_block:

```
ID   V38663 standard; DNA; 2807 BP.
AC   V38663;
DT   27-OCT-1998 (first entry)
DE   Rattus norvegicus SOCS1 gene.
KW   SOCS; suppressor of cytokine signalling; SOCS1 gene.
KW   autoimmune disease; diagnosis; cancer; treatment;
KW   cytokine mediated cellular responsiveness; hyperimmunity;
```

KW immunosuppression; allergies; hypertension; ss.
OS Rattus norvegicus.
FH Key   Location/Qualifiers
CDS   1739..2377
                /*tag= a
                /product= SOCS1 protein
FT
FT
PN WO9820023-A1.
PD 14-MAY-1998.
PF 31-OCT-1997; AU0729.
PF 14-FEB-1997; AU-005117.
PR 01-NOV-1996; AU-003384.
PA (HALL-) HALL INST MEDICAL RES WALTER & ELIZA.
PI Alexander WS, Hilton DJ, Nicholson SE,
PI Nicola NA, Richardson RT, Starr R, Viney EM, Willson TA;
DR WPI; 98-286854/25.
DR P-PSDB; W62617.
PS Claim 14; Page 117-118; 325pp; English.
PT The sequence is that of a gene encoding a suppressor of cytokine
PT signalling protein (SOCS). SOCS can be used to screen for naturally
PT occurring antibodies to SOCS, which may occur, e.g. in some autoimmune
CC diseases. Alternatively, specific antibodies can be used to
CC screen for SOCS, which is useful as a knowledge of SOCS levels
CC may be important for the diagnosis of certain cancers. Soluble
CC SOCS polypeptides can be used to treat disease, injury or
CC abnormality involving cytokine mediated cellular responsiveness,
CC e.g. hyperimmunity, immunosuppression, allergies and hypertension.
SQ Sequence 2807 BP;   906 C;   899 G;   495 T;

alignment_scores:
        Quality: 1073.00   Length: 212
        Ratio: 5.134   Gaps: 0
Percent Similarity: 98.585   Percent Identity: 96.226

alignment_block:
US-08-962-560A-4 x V38663

Align seq 1/1 to: V38663 from: 1 to: 2807

        1 MetValAlaAlaArgAsnGlnValAlaAlaAsnAlaIleSerProAlaAl 17
          ||||||||||||||||||||||||||||||||||||||||||||||||:||
     1739 ATGGTAGCAGTAACCAGGTGGAACCCGACCAAATGCGAATCNTCCCCGGCATC 1788

       17 aGluProArgArgArgSerGluProSerSerSerSerSerSerSerSerP 34
          ||||||||||||||||||||||||||||||||||||||||||||||||||
     1789 AGAGCCCCGACGGGCGGCCCAGAGCCATCCTCGTCCTCGTTCGTCGTCTCGC 1838

       34 roAlaAlaAlaProValArgProCysProAlaValProAlaProAla 50
          ||||||||||||||||||||||||||||||::|||||||||||||||||
     1839 CGGCGGCCCCCGGCCCCGGCGGGTCCCCGGCCCGTGGGTCCGGCGCCCCGGCT 1888

       51 ProGlyAspThrHisPheArgThrPheArgSerHisSerAspTyrArgAr 67
          |||||||||||||||||||||||||||||||||||||||||||||||||
     1889 CCGGGCGGACACTCACTCCCACCTTCCGGTCCCACTCTGATTACCGGCG 1938

       67 gIleThrArgThrSerAlaLeuLeuAspAlaCysGlyPheThrTrpGlyP 84
          |||||||||||||||||||||||||||||||||||||||||||||||||||
     1939 CATCACGCGGACACCAGCGCTCTCCTGGACGCCTGCCGCTTCTACTGGGAC 1988

       84 roLeuSerValHisGlyAlaHisGluArgLeuArgAlaGluProValGly 100
          |||||||||||||||||||||||||||||||||||||||||||||||||||
     1989 CCCCTGAGCGTGCATGGGGCGCACGAACGGCTGCGTTCCGAACCCGTGGGC 2038

      101 ThrPheLeuValArgAspSerArgGlnArgAsnCysPheGluAlaLeuSe 117
          |||||||||||||||||||||||||||||||||||||||||||||||||||
     2039 ACCTTCCTTGGTGCGCGACAGCGCCAGCGGAACTCCTTCTTCGGCCGTCAG 2088

      117 rValLysMetAlaSerGlyProThrSerIleArgValHisPheGlnAlaG 134
          |||||||||||||||||||||||||||||||||||||||||||||||||||

seq_name: N_Geneseq_36:V38662

seq_documentation_block:
ID V38662 standard; DNA; 1094 BP.
AC V38662;
DT 27-OCT-1998 (first entry)
DE Homo sapiens SOCS1 gene.
KW SOCS; suppressor of cytokine signalling; PCR primer;
KW autoimmune disease; diagnosis; cancer; treatment;
KW cytokine mediated cellular responsiveness; hyperimmunity;
KW immunosuppression; allergies; hypertension; ss.
OS Homo sapiens.
FH Key   Location/Qualifiers
CDS   24..659
                /*tag= a
                /product= SOCS1 protein
FT
FT
PN WO9820023-A1.
PD 14-MAY-1998.
PF 31-OCT-1997; AU0729.
PF 14-FEB-1997; AU-005117.
PR 01-NOV-1996; AU-003384.
PA (HALL-) HALL INST MEDICAL RES WALTER & ELIZA.
PI Alexander WS, Hilton DJ, Nicholson SE,
PI Nicola NA, Richardson RT, Starr R, Viney EM, Willson TA;
DR WPI; 98-286854/25.
DR P-PSDB; W62616.
PS Claim 14; Page 115-116; 325pp; English.
PT The sequence is that of a gene encoding a suppressor of cytokine
PT signalling protein (SOCS). SOCS can be used to screen for naturally
PT occurring antibodies to SOCS, which may occur, e.g. in some autoimmune
CC diseases. Alternatively, specific antibodies can be used to
CC screen for SOCS, which is useful as a knowledge of SOCS levels
CC may be important for the diagnosis of certain cancers. Soluble
CC SOCS polypeptides can be used to treat disease, injury or
CC abnormality involving cytokine mediated cellular responsiveness,
CC e.g. hyperimmunity, immunosuppression, allergies and hypertension.
SQ Sequence 1094 BP;   167 A;   381 C;   313 G;   233 T;

alignment_scores:
        Quality: 1053.50   Length: 212
        Ratio: 5.041   Gaps: 1
Percent Similarity: 98.585   Percent Identity: 95.283

alignment_block:
US-08-962-560A-4 x V38662

     2089 CGTGAAGAATGGCTTCGGGCCCCACGAGCATTCGTGGTGCACTTCCAGGCCG 2138
      134 lyArgPheHisLeuAspSpGlySerArgGluThrPheAspCysLeuPheGlu 150
          |||||||||||||||| |||::||||||||||||||||||||||||||||
     2139 GCCGCTTCCACCTTGCCAGCGGCCAACCGGCCGGAGACCTTCGACTGCCTTTGCGAG 2188
      151 LeuLeuGluHisTyrValAlaAlaLeuProArgArgMetLeuGlyAlaProLe 167
          ||||||||||||||||||||||||||||||||||||||||||||||||||
     2189 CTGCTGGAGCACTACGTGGCGGCGCTGCCGCGGCGCATGTTGGGGGCCCCACT 2238
      167 uArgGlnArgArgValArgProLeuGlnGluLeuCysArgGlnArgIlev 184
          ||||||||||||||||||||||||||||||||||||||||||||||||||
     2239 GCGCCAGCGCCGCGTGCGGCCCCTGCAGGAGCTGTGTCGCCAGCGCATCG 2288
      184 alAlaAlaValGlyArgGluAsnLeuAlaArgIleProLeuAsnProVal 200
          ||||||||||||||||||||||||||||||||||||||||||||||||||
     2289 TGGCCGCCGTGGGTCGGGAGAACCTGGCCAGGGATCCCCTCTTAACCCGTA 2338
      201 LeuArgAspTyrLeuSerSerPheProPheGlnIle 212
          |||||||||||||||||||||||||||||||||||||
     2339 CTCCGGTGACTACCTGAGTTCCTTCCCCTTCCAGATC 2374

Align seg 1/1 to: V38662 from: 1 to: 1094

```
   1 MetValAlaArgAsnGlnValAlaAlaAspAsnAlaIleSerProAlaAl 17
     ||||||||||::|::||||||||||||||||||||||||||||||||||
  24 ATGGTAGCACACAACCAGGTGGCAGCCGACAATGCAGTCTCCACAGCAGC 73

  17 aGluProArgArgSerGlnProSerSerSerSerSerSerSerSerSerP 34
     |||||||||::||||||| |||||||||||||||||||||||||||||
  74 AGAGCCCGACGGGCGGCCAGAACCT...TCCTCCTCTTCCTCCTCTCCGC 120

  34 roAlaAlaProValAlaProArgProCysProAlaValProAlaProAla 50
     |||||||:|::|||||||||||||||||||||||||||||||||||||||
 121 CCGGCGGCCCCCGCGCCCGCGGCCTGCCCGGCGGTCCCGGCGCCCGGCC 170

  51 ProGlyAspThrHisPheArgThrPheArgSerHisSerAspTyrArgAr 67
     ||||||||||||::|||:||||||||||||||||||||||||||||||
 171 CCCGGCGACACGCACTTCCCACATTCCGGTTCGCACGCCGATTACCGGCG 220

  67 gIleThrArgThrSerAlaLeuLeuAspAlaCysGlyPheTyrTrpGlyP 84
     ||||||||:||:||||||||::|||||||||||||||||||||||||||
 221 CATCACGCGCCAGCCGCCTCCTCGGACCCCTGCGGGATTCTACTCGGGGGC 270

  84 roLeuSerValHisGlyAlaHisGluArgLeuArgAlaGluProValGly 100
     ||||||||||||||||||||||||||||||||||||||||||||||||
 271 CCCTGAGCGGTGCACGGGCGCCACGAGCGGCTGCGCGCGGAGCCCGTGGGC 320

 101 ThrPheLeuValArgAspSerArgGlnArgAsnCysPhePheAlaLeuSe 117
     ||||||||||||||||||||||||||||||||||||||||||||||||
 321 ACCTTCCTGGTGCGCGACAGCCGCCAGCGAACTGCTTTTTCGCCCCTTAG 370

 117 rValLysMetAlaSerGlyProThrSerIleArgValHisPheGlnAlaG 134
     ||||||||||||||||||||||||||||||||||||||||||||||||
 371 CGTGAAGATGGCCTCGGGACCCACGAGCATCCGCGTGCACTTTCAGGCCG 420

 134 lyArgPheHisLeuAspGlySerArgGluThrProAspCysLeuPheGlu 150
     ||||||||||||||||||||||||||||||||||||||||||||||||
 421 GGCCGCTTTCACCTGGATGGCAGCCGCGAGACTCCGGACTGCCTCTTCGAG 470

 151 LeuLeuGluHisTyrValAlaAlaProArgMetLeuGlyAlaProIle 167
     ||||||||||||||||||||||||||||||||||||||||||||||||
 471 CTGCTCGGAGCACTAGGTGGGGCGGCGCATCCTGGGGGCGCCCCGCT 520

 167 uArgGlnArgGlyValArgProLeuGlnGluLeuCysArgGlnArgIleV 184
     ||||||||||||||||||||||||||||||||||||||||||||||||
 521 GCGCCAGCCGCCGGGCCTGCGCCCGGCGGAGCTGTGCCGCCAGCGCATCG 570

 184 aLAlaAlaValGlyArgGluAsnLeuAlaArgIleProLeuAsnProVal 200
     |||||:::||||||||||||||||||||||||||||||||||||||||
 571 TGGCCACCGTGGGCCGCGAGAACCTGGCTCGCATCCCCTTCAACCCGGTC 620

 201 LeuArgAspThrTyrLeuSerSerSerPhePheProGlnIle 212
     |||||||||||||||||||||||||||||||||||||||||
 621 CTCCGCGACTACCTGAGCTCCTTCCCTTCCCAGATT 656
```

seq_documentation_block:

seq_name: N_Geneseq_36:T43380

alignment_scores:

```
             Quality:   249.50        Length:      297
             Ratio:     1.835         Gaps:         10
Percent Similarity: 45.117   Percent Identity: 27.946
```

alignment_block:
US-08-962-560a-4 x T43380

Align seg 1/1 to: T43380 from: 1 to: 1960

```
  14 SerProAlaAlaGluProArgArgSerGlnProSerSerSerSerSe 30
     ::|||||||||||:|||  |||  ||||| :::||||
  10 GCCCCGGGAGCCTACCCAGCACGCGCCCGCGGCCCACTGGTTCCCTCCAG 59

  30 rSerSerSerProAla..AlaProValArg.................. 39
     |||||||||||||  :::||:|::||||
  60 CCGCCCCGTCCCAGCCGAGTCCCCACTCCGGAGTCGCCCGGTGCCGCGGGG 109

  40 ..............ProArgProCysProAlaVal........ 46
                  ||||||||||||
 110 ACATGGTCCTCTGCGTTCCAGGGACCTTCGTTCCTTTGCTGGCGTTGGAGCGG 159

  46 ............................ProAla.............. 48
                              ||||||
 210 CATGCAGCCCTTGCCTGGGGGCCGTTCCTCGAGGAGGTGGCAGAGGGTA 259

  49 ..ProAlaProGlyAspThrHisPheArgThrPheArgSerHisSerAsp 64
       |||||| :::||:||:||::||||  :::  ::||:
 260 CCCCAGCCCAGACAGAGAGTGAGCCAAAAGGTGCTGGACCAGAGGAGGAT 309

  65 TyrArgArgIleThrArgThrSerAlaLeuLeuAspAlaCysGlyPheTy 81
     ||:||:||||||||||  :::||:
 310 CTGCGTGTGCAATAGCCAAGACCTTCTCCTACCTTCGGGAATCTGGCTGGTA 359

  81 rTrpGlyProLeuSerValHisGlyAlaHisGluArgLeuArgAlaGluP 98
     ||||| |:::::::|||  ||||  ||:::|||
 360 TTGGGGTTCCATTACGGCCAGCGAGCCCGACAACACCTCGCAGAAGATGC 409

  98 roValGlyThrPheLeuValArgAspSerArgGlnArgAsnCysPhePhe 114
       ||||  ||::||||||||:::||:||
 410 CAGAAGGCACGTTCTTAGTACGTGACACAGCACCCCAGCTACCGTTC 459

 115 AlaLeuSerValLysMetAlaSerGlyProThrSerIleArgValHisph 131
     ::|||||||||||   :::||||  |||:::::||||::||||||
 460 ACGCTGTCAGTGAAAACCACTCGTGCCCCACCAATGTACGCAGTGAGTA 509

 131 eGlnAlaGlyArgPheHisLeuAspGly................SerArgGlu 143
     ::|||||  |||::::::::::                :::||||
```

seq_name: N_Geneseq_36:V38660

```
510 TGCCGACTCCAGCTTCCGTCTGGACTCCAACTGCTTGTCCAGGCACGCA 559
        ||||::::||||||||||
144 ......ThrPheAspCysLeuPheGluLeuLeuGluHisTyrValAla... 157
560 TCCTGGCCTTTCCGGATGTGGTCAGCCTTGTCAGCACTATGTGGCCTCC 609
        ::||||||||||||||||||||||||||
158 ...........................AlaProArgArgMe 162
610 TGCACTGGTTGGATACCCGGAAGGCACGACCCCGAATCCTCCACCCGGC 659
        |||||||
162 tLeuGlyAlaPro......................AlaProArgArgMe 166
660 CCTGCCTATGCCTAAGGAGGATGCCCTAGTGACCCAGCACTGCCTGCTC 709
        |:||||||||||||||||
167 ................................LeuArg 168
710 CTCCACCAGCCACTGCTGTACACCTAAAACTGGTGCAGCCCTTTGTACGC 759
        ::||||
167 ...................................SerSerSerSerSerPro 184
760 AGAAGAAGTGGCCCCGCCAGCTGCTGCAACACGTGGCGCCCTTGTCATCAACG 809
        |||||||||::||||||||||||::
169 GlnArgArgValArgProLeuGlnGlnValCysArg.........GlnAr 182
810 TCTGGTGGCCGACGTGGAC.................TGCCTGCCACTGCCCC 844
        |:::||:::||||
182 gIleValAlaValGlyArgGluAsnLeuAlaArgIleProLeuAsnP 199
845 GGGGCCATGCCGCGACTACCTCCGACAGTACCCCTCCAGCTC 885
        ||||||||    :::|||||||||||||||::
199 roValLeuArgAspTyrLeuSerSerPheProPheGlnIle 212
```

seq_documentation_block:

alignment_scores:
      Quality: 234.50   Length: 248
      Ratio: 1.777   Gaps: 9
      Percent Similarity: 53.226   Percent Identity: 29.032

alignment_block:
      US-08-962-560A-4:

Align seg 1/1   to: V38660   from: 1   to: 1121
                       x V38660

```
14  SerProAlaAlaGluProArgArgArgSerGluProSerSer..... 28
        |:||||||||      ||    :
85  TCCCCTCCCCCCTCCCGCCACCATTTCGGACACCTCGTTTTG 134
29  ...............SerSerSerSerSerProAlaAlaProValA 39
        ::||||:::    ||||||:::||
135 GGGTACCCTGACTTCCAGGCACGCAGGAGGTTCCACTGGCCCCAGCTC 184
39  rgProArgProCysProAlaValProAlaProAlaPro......... 51
        |||  :::              ::|
185 GGGCGACCAGCGTGTTGACGTGTTGACTCATCCCATGACCCTGCGG 234
52  .......GlyAspThrHisPheArgThrPheArgSerHis.. 62
        |:||||::    :::::||||
235 TGCCTGGAGCCCTCCGGGAATGGAGCGGGACAGGACG...CGGAGCCAGTG 281
63  ...........................SerAspTyrArgArgIle 69
        :::                  ::|||
282 GGGGACCGCGGGGGTTGCCGGAGGACAGTCCCCAGGGCGCGCTCTGG 331
69  hrArgThrSerAlaLeuLeuAspAlaCysGlyPheTyrTrpGlyProLeu 85
        ||||:|      |||||||||||||:
332 CGAAAGCCCTGCGCCGAGCTCAAACAGGATGGTACGGGAGGAAGTATG 381
86  SerValHisGlyAlaHisGluArgLeuArgLeuGluProValGlyThrPh 102
        ||||    |||::::||||      ||||||
382 ACTGTTAATGAAGCCAAAGAGAAAATTAAAAGAGGCTCCCAGAAGGAACTTT 431
102 eLeuValArgAspSerArgGlnArgAsnCysPhePheAlaLeuSerVall 119
        |||:|      ||||||::::      :::|||||
432 CTTGAATTAGAGATAGTTCCCATTCAGACACTACCTAACTATATCCGTTA 481
119 ysMetAlaSerGlyProThrSerIleArgValHisPheGlnIleGlyArg 135
        ||      :||||||::||||||::::||        |||:||
482 AGACGTCAGGTGGCACCGGACTAACCTGCGGATTGAGTACCAAGAGGAAA 531
136 PheHisLeuAsp...................GlySerArgGlyThrPheAs 146
        ||      ||||::
532 TTCAGATTGGATTCTAATCAATATGTGTCAAGTCCAAGCTTAAACAGTTTGA 581
146 pCysLeuPheGluLeuLeuGluHisTyrVal.................. 156
        :::                  ||||||
582 CAGTGTGGTTCATCTGATTGAACTACTATGTCCAGATGTGCAAGGATAAAC 631
157 ....................AlaAlaProArgArg............... 164
        ||||||||:::
632 GGACAGGCCCAGAAGCCCAGGAACTGGACTGGTTCACCGTACCTGACC 681
165 AlaProLeuArgGlnArgArgValArgProLeuGlnGlnLeuCysArgGl 181
        |||||:::                  ||||||:::|||
682 AAAACCTCTG...TATACACATGAGCACCCACTCCTGCAGCATTTCTGTCGACT 728
181 nArgIleValAlaValGlyArgGluAsnLeuAlaArgIleProLeuLeuA 198
        |||                  :::||||||
729 CGCCAATTAACAAAATGTACCGGT.......ACGATCTGGGGACTGCCTTTAC 772
198 snProValLeuArgAspTyrLeuSerSerPheProPheGlnIle 212
        |||:|||||||      ||||||||||||||:::
773 CAACAAGACAACAAAAGAATTACTTGGAAGAATAAAATTCCAGGTA 816
```

seq_name: N_Geneseq_36:V69307

seq_documentation_block:
ID   V69307   standard; cDNA; 2342 BP.
AC   V69307;
DT   01-FEB-1999   (first entry)
DE   Human EPRG1 cDNA #1.
KW   EPRG1; EPO primary response gene 1; diagnosis; gene therapy; immunity;
KW   disease; vaccine; inoculate; antibody; T cell; anaemia; polycythemia;
KW   cancer; neutropenia; AIDS; diabetes, myelosuppression; allergy; asthma;
KW   autoimmune disease; inflammatory disease; chromosome mapping; human; ss.
OS   Homo sapiens.
PN   EP-877030-A2.
PD   11-NOV-1998.
PF   07-MAY-1998; 303597.
PR   01-MAY-1998; US-071342.
PR   07-MAY-1997; US-045890.
PA   (SMIK) SMITHKLINE BECKMAN CORP.
PI   Dillon S, Lord K;
WPI 98-570499/49.
DR   P-PSDB; W82504.
PT   New EPO primary response gene polypeptides and polynucleotides -
PT   useful as diagnostic reagents and for prevention and treatment of
PT   cancer and autoimmune and inflammatory diseases
PS   Claim 14; Page 18-19; 25pp; English.
CC   This sequence encodes a novel human EPO primary response gene 1 (EPRG1)
CC   polypeptide. EPRG1 polypeptides and polynucleotides are useful for
CC   diagnosing a disease or susceptibility to a disease by detecting
CC   mutations in the EPRG1 gene using probes containing the EPRG1 nucleotide
CC   sequence, or determining EPRG1 polypeptide or mRNA expression levels
CC   (antagonist eg EPRG1 ligand, receptor or substrate) EPRG1 activity, in
CC   EPRG1 polypeptides can be used to screen for agonists and antagonists
CC   which bind the EPRG1 polypeptide by measuring resulting mRNA levels with
CC   ELISA. These can be used in treatment to activate (agonist) or inhibit
CC   addition to direct administration of antisense sequences to prevent
CC   expression, or EPRG1 polypeptides can be used to treat conditions associated with
CC   a lack of EPRG1 protein. Gene therapy may also be used to affect
CC   endogenous EPRG1 polypeptide production. EPRG1 antibodies are useful for
CC   isolating EPRG1 clones or purifying the polypeptides by affinity
CC   chromatography. EPRG1 polypeptides can be administered directly or as a
CC   vaccine to inoculate against disease by inducing an antibody and T-cell
CC   response. Diseases diagnosed, prevented or treated include anaemia,
CC   polycythemia, cancer, neutropenia, AIDS, drug-induced anaemia, diabetes,
CC   myelosuppression, autoimmune diseases, rheumatoid arthritis and multiple
CC   sclerosis, and inflammatory diseases, including asthma and allergies. The
CC   EPRG1 polypeptide is also useful for mapping the gene to a chromosome,
CC   allowing gene inheritance to be studied through linkage analysis. The
CC   3'-UTR segment of EPRG1 RNA may be useful to screen for agents which
CC   modulate RNA stability and turnover rate.
SQ   Sequence 2342 BP; 495 A; 685 C; 655 G; 506 T;

alignment_scores:
            Quality: 234.50   Length: 231
              Ratio: 1.804   Gaps: 9
Percent Similarity: 56.277   Percent Identity: 29.870

alignment_block:
US-08-962-560A-4 x V69307

Align seg 1/1 to: V69307   from: 1 to: 2342

```
30  SerSerSerProAlaAlaProValaArgProArgProCysProAlava  46
    |||::||     ::||||| :||   |||   ||||:||:
7   TCCACGCnGGCTCCGTGCGCCAnGGTCACCCAGCAAGTTCCCGCCGC  56

46  IproAlaProAlaProGlyAspThrHisPheArg........ThrPheArg  61
    :  ||||||||:||||||||||||:           ||||| |
57  CGGGATGAGCCGCGCCCCCTGGACACCAGCCTGCGCCTCAAGACCTTCAGCT  106

61  erHisSerAspTyrArgArgIleThrArgThrSerAlaLeuLeuAspAla  77
    ||     ||||:||||:||||::|||||||
107 CCAAGAGGCGAGTACCAGCGTGGTGGTGAACGCACTGCGCAAGCTGCAGGAG  156
```

---

seq_documentation_block:
ID   V69309   standard; cDNA; 2342 BP.
AC   V69309;
DT   01-FEB-1999   (first entry)
DE   Human EPRG1 cDNA derived from expressed sequence tags, EST's.
KW   EPRG1; EPO primary response gene 1; diagnosis; gene therapy; immunity;
KW   disease; vaccine; inoculate; antibody; T cell; anaemia; polycythemia;
KW   cancer; neutropenia; AIDS; diabetes, myelosuppression; allergy; asthma;
KW   autoimmune disease; inflammatory disease; chromosome mapping; human;
KW   expressed sequence tag; EST; ss.
OS   Homo sapiens.
PN   EP-877030-A2.
PD   11-NOV-1998.
PF   07-MAY-1998; 303597.
PR   01-MAY-1998; US-071342.
PR   07-MAY-1997; US-045890.
PA   (SMIK) SMITHKLINE BECKMAN CORP.
PI   Dillon S, Lord K;
WPI 98-570499/49.
PT   New EPO primary response gene polypeptides and polynucleotides -
PT   useful as diagnostic reagents and for prevention and treatment of
PT   cancer and autoimmune and inflammatory diseases
PS   Claim 13; Page 22-23; 25pp; English.
CC   This sequence encodes a novel human EPO primary response gene 1 (EPRG1)
CC   polypeptide derived from expressed sequence tags (EST's). EPRG1
CC   polypeptides and polynucleotides are useful for diagnosing a disease or
CC   susceptibility to a disease by detecting mutations in the EPRG1 gene

seq_name: N_Geneseq_36:v69309

```
78  CysGlyPheTyrTrpGlyProLeuSerValHisGlyAlaHisGluArgIle  94
    |||||||||||||||:::       :::::            ||
157 AGCGGCTTCTACTGGGAGGGCCAGTAGGGCGGCGGGGGCGAGGCGAACCTGCTCCT  206

94  uArgAlaGluProValGlyThrPheLeuValArgArgSerArgGlnArgA  111
    :||||||||:||||||||||||:||:|||    |||::          ||
207 CAGTGCCGGAGCCCGGGCGCACCTTTCTGGTGCGCCGCAGCCGCGACAGCTCGGACCAGC  256

111 snCysPhePheAlaLeuSerValLysMetAlaSerGlyProThrSerIle  127
    |||||||::||:||||||||||||:|||||:|||||||
257 GCCACTTCTTCACGCTCAGCGTCAAGACCCAGTGTGGGACCAGAGACCTG  306

128 ArgValHisPheGlnAlaGlyArgPheHisSerLeuAspPheGlySer...ArgG  143
    :||||||:||||||:||     |||    |||::
307 CGCATCCACGTGTGAGGGGCGGCAGCTTCACTTCTCTGCAGACGGATCCCGGAG  356

143 uThr...........PheAspCysLeuPheGluLeueuGluHisr  155
    :|||                      |||||||||::|||||||||
357 CACGCAGCCGCGGTGCCCGCGGCTTCGACTGCCTGCTCAAGCTGGTGCACCACT  406

155 yrVal...........  156
    ||::
407 ACATGCCGCCCCCTG  456

157                                                     AlaAl  158
    :::
457 TCCTCCGAGGTGCCGAGCAGCCGGTCTCCCAGCCACTCCCTGGGAGTCC  506

158 aProArgArg..........................  M  162
    ::||||||                                   :
507 CCCCAGAGAGCCTATTACATCTACTCCGGGGGCGGAGAAGAATCCCCTGG  556

162 etLeuGlyAlaProLeuArgGlnArgArgValArgProLeuGlnGluLeu  178
    :::||||||   |||||| :|||:||         |||||||||:::|||
557 TGTTGAGCCGCGGCCCCTC...TCCTCCAACGTGGCCACTCTTCAGCATCTC  603

179 CysArgGlnArgIleValAlaAlaAlaVal...GlyArgGluAsnLeuAlaAr  194
    :||||||| ::  |||||::      ::         ||||||:::::::
604 TGTCGGCAGAGACCGTCAACGGCGCCACCTGACTCCTATGAGAAAGTCACCCA  653

194 gIleProLeuAsnProValLeuuArgAspTyrLeuSerSerPhe  208
    ||         |||:||::||     ||::::::::
654 GCTGCCG...GGGCCCATt...CGGGAGTTCCTGGACCAGTAC  690
```

```
CC  using probes containing the EPRG1 nucleotide sequence, or determining
CC  EPRG1 polypeptide or mRNA expression levels EPRG1 polypeptides can be
CC  used to screen for agonists and antagonists which bind the EPRG1
CC  polypeptide by measuring resulting mRNA levels with ELISA. These can be
CC  used in treatment to activate (agonist) or inhibit (antagonist eg EPRG1
CC  ligand, receptor or substrate) EPRG1 activity, in addition to direct
CC  administration of antisense sequences to prevent expression, or EPRG1
CC  polypeptides to treat conditions associated with a lack of EPRG1 protein.
CC  Gene therapy may also be used to affect endogenous EPRG1 polypeptide
CC  production. EPRG1 antibodies are useful for inducing an immune response
CC  to immunise and prevent diseases, and for isolating EPRG1 clones or
CC  purifying the polypeptides by affinity chromatography. EPRG1 polypeptides
CC  can be administered directly or as a vaccine to inoculate against disease
CC  by inducing an antibody and T-cell response. Diseases diagnosed,
CC  prevented or treated include anaemia, polycythemia, cancer, neutropenia,
CC  AIDS, drug-induced anaemia, diabetes, myelosuppression, autoimmune
CC  diseases, rheumatoid arthritis and multiple sclerosis, and inflammatory
CC  diseases, including asthma and allergies. The EPRG1 polypeptide is also
CC  useful for mapping the gene to a chromosome, allowing gene inheritance to
CC  be studied through linkage analysis. The 3'-UTR segment of EPRG1 RNA may
CC  be useful to screen for agents which modulate RNA stability and turnover
CC  rate.
SQ  Sequence  2342 BP;  495 A;  685 C;  655 G;  506 T;

alignment_scores:
         Quality:  234.50
         Ratio:    1.804
Percent Similarity:  56.277    Percent Identity:  29.870

alignment_block:
US-08-962-560A-4 x V69309

Align seg 1/1  to: V69309    from: 1  to: 2342      Length:   231
                                                    Gaps:       9

30  SerSerSerSerProAlaAlaProValArgProArgProCysProAlaVa  46
    |||||:::||||||:::||  ||
7   TCCACGCTGGCTCCGTGCCCCATGGTCACCCACACAGCAAGTTTCCGCCCC  56

46  1ProAlaProAlaProGlyAspThrHisPheArg.......ThrPheArgs  61
    |||  |||||  :::|||              ||||||  |
57  CGGGATGAGCCGCCCCCCTGGACACCAGCCTGCGCCTCAAGACCTTCAGCT  106

61  erHisSerAspTyrArgArgIleThrArgThrSerAlaLeuLeuAspAla  77
    |:|||||:::||||||||:::|:::||||||          |||||:
107 CCAAGAGCGAGTACCAGCGTGGTGGAAGCGAGTGCGGAAGCTGCAGGAG   156

78  CysGlyPheTyrTrpGlyProLeuSerValHisGlyAlaHisGluArgle  94
    :||||||||||||:||||||  |||||||||||:::|||:::||:
157 AGGGGCTTCTACTGGAGGCCAGTGAACCGGCGGCGAGGCGAACCTGCTGCT  206

94  uArgAlaGluProValGlyThrPheLeuValArgAspSerArgGlnArga 111
    |  ||||||||:::|||||||||:||||||:::||||||||||||
207 CAGTGCGCCAGCCGCGGCCACCTTTCGATCCCCGACAGCTCGGACCAGAGC  256

111 snCysPhePheAlaLeuSerValLysMetAlaSerGlyProThrSerIle  127
    ::  |||||:::||||||||:::||||||||||:::||||:::
257 GCCACTTCTTCACGCTCAGCGTCAAGACCCAGTCTGGGACCAAGAACCTG  306

128 ArgValHisPheGlnlnAlaGlyArgPheHisLeuAspGlySer....ArgGl 143
    |||||||||:::::||||  |||  ||||||||||||||||||  |||::
307 CGGATCCAGTGTGAGGGGGGCAGCTTCTCTCTGGCAGAGGGATCCCCGGAG  356

143 uThr...........PheAspCysSerLeuPheGluLeuLeuGluHisI  155
    :|||           ||||||||||||:::|||||||:::|||||
357 CACGCAGCCCGTGCCCCCCTTCGACTGCGTCGTCAAGCTGGTCCACCACT  406

155 yrVal..........  156
    |:::|
407 ACATGCCGCCCCCCTGGAGCCCCCTCCCCCTGCCACCTACTGAACCC  456

157 ..........  157
    :::
                                          .AlaAl 158
                                          :::
```

```
457 TCCTCCGAGGTGCCCGAGCAGCCCGTGCCCAGCCGCCGACTCCCTGGGAGTGCC 506
    ...................................................M 162
158 aProArgArg..............  162
    |||||
507 CCCAGAAGAGGCCTATTACATCTACTCCGGGGGCGGCGGAGAAGAATCCCCTGG 556
                                                          :
162 etLeuGlyAlaProLeuArgGlnArgArgValArgProLeuGlnGluLeu 178
    :::|||:::  |||||:  :::  ||||||:::  ||||||:::  ||||
557 TGTTGAGCCGGCCGCCCTC...TCCTCCACGTGCCACTCTTCAGCATCTC 603

179 CysArgGlnArgIleValAlaLeuAlaVal...GlyArgGluAsnLeuArg 194
    |||:::|||:::  :::  |||:::  ||||||  :::||:::|||:::
604 TGTCGGAAGACCGTCAACCGCCACCTGGACTCCTATGAAAAGTCACCA 653

194 gIleProLeuAsnProValLeuArgAspTyrLeuSerSerPhe 208
    :::|||:::|||:::  |||:::|||:::||||||||||:::
654 GCTGCCG...GGGCCCATT...CGGGAGTTCCTGGACCAGTAC 690
```

```
seq_name: N_Geneseq_36:T96002

seq_documentation_block:
ID  T96002 standard; cDNA; 1374 BP.
AC  T96002;
DT  07-JUL-1998  (first entry)
DE  Human cytokine-inducible SH2-containing (CIS) gene.
KW  Human cytokine-inducible SH2-containing; CIS; erythroprotein; EPO;
KW  tyrosine-phosphorylated interleukin; (IL)-3; signal transduction;
KW  inhibition; anaemia; ss.
OS  Homo sapiens.
FH  Key            Location/Qualifiers
FT  CDS            71..848
FT                 /tag= a
FT                 /product= "Human CIS protein"

PN  WO9744347-A1.
PD  27-NOV-1997.
PF  21-MAY-1996;  WO-U07477.
PR  21-MAY-1996;  WO-U07477.
PA  (HARD ) HARVARD COLLEGE.
PA  (SMIK ) SMITHKLINE BEECHAM CORP.
PI  Dunnington DJ, Frantz JD, Shoelson SE;
DR  WPI; 98-018425/02.
DR  P-PSDB; W38319.
PT  Human cytokine-inducible SH2-containing protein and related DNA -
PT  useful for diagnosis of modulators for treatment of, e.g. anaemia
PS  Claim 1; Pages 34-35; 52pp; English.
CC  This sequence encodes the human cytokine-inducible SH2-containing
CC  (CIS) protein which binds the human cytokine-inducible interleukin
CC  (IL)-3 or erythroprotein (EPO) receptors. When CIS is over expressed
CC  signal transduction through the receptors is inhibited. Inactivation
CC  of CIS may enhance signalling through (IL)-3 and EPO receptors.
CC  Specific inhibitors of CIS may be useful in the treatment of anaemia.
CC  Labelled CIS or its functional derivatives can be used in binding
CC  assays to determine modulators of CIS activity. Conditions associated
CC  with CIS protein deficiency can be diagnosed by assaying for the
CC  presence of the CIS gene. It can be used to treat conditions related
CC  to insufficient CIS protein function.
SQ  Sequence  1374 BP;  264 A;  446 C;  379 G;  285 T;

alignment_scores:
         Quality:  234.00
         Ratio:    1.759
Percent Similarity:  46.503    Percent Identity:  26.573

alignment_block:
US-08-962-560A-4 x T96002

Align seg 1/1  to: T96002  from: 1  to: 1374      Length:   286
                                                  Gaps:      10

19  ProArgArgSerGluProSerSerSerSerSerSerSerSerProA1 35
    |||:::|||:::  :::|||
18  CCGAAGCGACAGCCGGATCCT............GCTCCCACTCCGGA 52
```

```
seq_name: N_Geneseq_36:V38661
seq_documentation_block:
ID   V38661 standard; DNA; 2187 BP.
AC   V38661;

DT   27-OCT-1998 (first entry)
DE   Mus musculus SOCS3 gene.
KW   SOCS; suppressor of cytokine signalling; SOCS3 gene;
KW   autoimmune disease; diagnosis; cancer; treatment;
KW   cytokine mediated cellular responsiveness; hyperimmunity;
KW   immunosuppression; allergies; hypertension; ss.
OS   Mus musculus.
FH   Key              Location/Qualifiers
FT   CDS              18..695
FT                    /*tag= a
FT                    /product= SOCS3 protein
PN   WO9820023-A1.
PD   14-MAY-1998.
PF   31-OCT-1997; AU0729.
PR   14-FEB-1997; AU-005117.
PR   01-NOV-1996; AU-003384.
PA   (HALL-) HALL INST MEDICAL RES WALTER & ELIZA.
PI   Alexander WS, Hilton DJ, Metcalf D, Nicholson SE,
PI   Nicola NA, Richardson RT, Starr R, Viney EM, Willson TA;
WPI  98-286854/25.
DR   P-PSDB; W62615.
DR   P-PSDB; W62615.
PS   Claim 14; Page 113-114; 325pp; English.
CC   The sequence is that of a gene encoding a suppressor of cytokine
CC   signalling protein (SOCS). SOCS can be used to screen for naturally
CC   occurring antibodies to SOCS, which may occur, e.g. in some autoimmune
CC   diseases. Alternatively, specific antibodies can be used to
CC   screen for SOCS, which is useful as a knowledge of SOCS levels
CC   may be important for the diagnosis of certain cancers. Soluble
CC   SOCS polypeptides can be used to treat disease, injury or
CC   abnormality involving cytokine mediated cellular responsiveness,
CC   e.g. hyperimmunity, immunosuppression, allergies and hypertension.
SQ   Sequence 2187 BP;                       632 C;       600 G;       454 T;
                                             501 A;

alignment_scores:
                Quality:    229.50        Length:    228
                Ratio:      1.779         Gaps:        9
     Percent Similarity:    56.579        Percent Identity:  29.386

alignment_block:
US-08-962-560A-4 x V38661

Align seq 1/1   to: V38661   from: 1   to: 2187
```

```
33  SerProAlaAlaProValArgProArgProCysProAlaValProAlaPr 49
    :::||||||::|||||:|||         ||
6   GGTCCGGTGCGCCAATGGTCACCCACAGCAAGTTTCCCGCGCCGGATGAG 55

49  oAlaProGlyAspThrHisPheArg.........ThrPheArgSerHisSerA 64
    |||   |||:||||:|:        |||:||
56  CCCGCCCCTGGACACCAGCCTCGGCCCTCAAGACCTTCAGCTCCAAAGCG 105

64  spTyrArgArgIleThrArgThrSerAlaLeuLeuAspAlaCysGlyPhe 80
    :|||:||    ||||||:|:|:|||       ||| ||||
106 AGTACCAGCTGGTGGTCGAACGCCGTGCCCAAGCTGCGAGGAGAGCGGATTC 155

81  TyrTrpGlyProLeuSerValHisGlyArgLeuArgArgValHis 97
    |||||:||||:|:|:|||:         ||| |||||
156 TACTGGAGCCGCCGTGACCGGCGCGGACCAAGCTGCTGCGCGCCA 205

97  uProValGlyThrPheLeuValArgAspSerArgGlnArgAsnCysHisPhe 114
    |:|||:      :::|||||      :||| ||||:
206 GCCCGCGGGCACCTTTCTTATCCGACAGCCGACAGGCCACTTCT 255

114 heAlaLeuSerValLysMetAlaSerGlyProThrSerIleArgValHis 130
    :|||:|::||||||||||||         |||||
256 TCACGTTGAGCGTCAAGACCCAGTCGGGGACGTCGGGGACCAAGAACCATCCAG 305

131 PheGlnAlaGlyArgPheHisLeuLeuAspLySer...ArgGluThr...... 144
```

```
35  aAlaProValArg........................ProArgProC 43
    ::|||:|||:|||
53  GTCGCGCGCTGCGGCGGAGACATGGTCTTCGGTTCAGGGACTCGTCCTT 102

43  ySProAlaVal........................ 46
    |||||
103 TGCTGCGTGTGGAGCGGACTGGGCAGCGGGGCCCGTGGGCCCGTCCCTG 152

47  ....................ProAla............. 48

153 GAACTGTCCAACCCAGTCATGCAGCCCTTGCCTGCTCTGGGGCCTTCCTCGA 202

49  ....................ProAlaProGlyAspThrHisPheArgThrP 59
    |||||           :::|||:|
203 GGAGGTGGGCAGAGGGTACCCAGCCACAGACAGAGTGAGCCAAAGTGC 252

59  heArgSerHisSerAspThrArgArgIleThrArgThrSerAlaLeuLeu 75
    :::|||:| |:||||:|:||||       |||  |||
253 TGGACCCAGAGGAGGAGGATCTGTGCATAGCCAAGACCTTCTCCTACCTT 302

76  AspAlaCysGlyPheTyrTrpGlyProLeuSerValHisGlyArgLeuArgG1 92
    :::|||:::  |||||||||:||||:|||:||:||:lyAlaHisGly
303 CGGGAAACTGGCTGGTTCCATTACGGCCACGGAGGCCCGACA 352

92  uArgLeuArgArgLeuGluProValGlyThrPheLeuValArgAspSerArgg 109
    ::|||:::|            |||||||||||||||||||
353 ACACCTGCAGAAGATGCCAGAAGGCACGTTCTTAGTACGTGACAGCAGACG 402

109 lnArgAsnCysHisPheheAlaLeuSerValLysMetAlaSerGlyProThr 125
    ::     ||:||||||||||||||||||||||||||||||||||||| |||
403 ACCCAACTACCTGTTCACGTTGAGCGTCAAGATGGCAAGCGGCCCCACC 452

126 SerIleArgValHisPheGlnAlaGlyArgPheHisLeuLeuAspAspGly 140
    :::||||||||||:||:|||:|||:|||||::
453 AATGTACGCATTGAGTATGCCGGACTCCAGCCTTCCGTGTCTGGACTCCAACTG 502

141 ....SerArgGlu.........ThrPheAspCysSerPheGluLeuLeuLeu 153
    :::|||
503 CTTGTCCAGGCCACGCAATCCTGGCCTTTCCGGATGTGGTCAGTCTTGTGC 552

153 luHisTyrValAla................LeuArgGlnArgArgValArgProLeuGlnGluCys 179
    ::|||:|||:|||
553 AGCACTATGTGGCCTTCCTGACTGCTGTATACCCGAAGCGACAGAGCCCGGAT 602

158 ...AlaProArgMetLeuGlyAlaPro.......................... 166
    |:||||| |||
603 CCTGCTCCACCCCGGTGTTCCTGCTATGCCTAAGGAGGATGCGGCCTAGTGA 652

166 .........................LeuArgGlnArgArgValArgProLeuGlnGluCys 179

653 CCCAGACACTGCCTCCTCACCCAGCCACCACTGCTGTACACCTAAAAACTGG 702

167 .................................................. 166
703 TGCAGCCCTTTGCTACGGCAGAAGCAGTGCCCCGCAGCCGCCAACACCTGC 752

180 Arg.................GlnArgIleValAlaAlaValGlyArgGluAsnLeuVal 193
    |||                 :::|||::||||||
753 CGCCTTGTCATCAAACCGTTCTGGTTGGCCGACAGTGGAC 788

193 aArgIleProLeuAsnProValLeuArgAspTyrLeuSerSerPheProP 210
    :|||                     |::|||
789 .TGCCTGCCACTGCCCCGGCGCGCCAACTACCTCCGACAGTACCCCT 837

210 heGlnIle 212
    ||:|||
838 TCCAGCTC 845
```

```
306 TGTGAGGGGGGGCAGCTTTTCGCTCGCTGCCAGAGTGACCCCCGAAGCACGCAGCC
      :::::|||    |||   |||::|::::::::|||   ||||::::::::|||
145 ........PheAspCysLeuPheGluLeuLeuGluHisTyrVal.... 156
                   ||||||||||||||||||::||||:::||:::
356 AGTTCCCGCGCTTCGACTGTGTACTCAAGCTGGTGCCACCACTACATGCCGC 405

156 ........ 156
      |||||||||||||||||||||||||||||||::||
406 CTCCAGGGACCCCCTCCTTTTCTTTGCCACCCACGGAACCTCGTCCGAA 455

157 ........AlaAlaProArgArg 161
      ::::||:||:||||
456 GTTCCGGAGCAGCCACCTGCCCCAGGCACTCCCCGGGAGTACCCCAAGAG 505

161 g.... MetLeuGlyA 165
      ::|||||:::
506 AGCTTACTACACATCTATTCTGGGGGGGGCGAGAAGATTCCGCTGGGTACTGACCC 555

165 laProLeuArgGlnArgArgValArgProLeuGlnGluLeuCysArgGln 181
      ||||| :|||:||||||||:|||||||:|
556 GACCTCTC..TCCTCCAACGTGGCCCACCCTCCAGCATCTTTGTCGGGAAG 602

182 ArgIleValAlaAlaVal...GlyArgGluAsnLeuAlaArgIleProLe 197
      :|| ::| |||||||||||||::::|||
603 ACTGTCAACGGCCACCTGGACTCCTATGAGAAAGTGACCCGCTGCCT.. 650

197 uAsnProValLeuArgAspTyrLeuSerSerPhe 208
      ::::||||:||:|||||||||:::
651 .GGACCCATT...CGGGAGTTCCTGGATCAGTAT 680
```

seq_name: N_Geneseq_36:V34188

seq_documentation_block:
```
ID   V34188 standard; DNA; 2378 BP.
AC   V34188;
DT   28-JAN-1999 (first entry)
DE   Human secreted protein gene 35 clone HTXAK60.
KW   Human; secreted protein; fusion protein; gene therapy; protein therapy;
KW   diagnosis; tissue; cancer; tumour; neurodegenerative disorder; leukaemia;
KW   development abnormality; foetal deficiency; blood; allergy; renal; ds;
KW   immune system; asthma; lymphocyte disease; brain; hepatic; lymphoma;
KW   inflammation; ischaemic shock; Alzheimer's disease; restenosis; AIDS;
KW   cognitive disorder; schizophrenia; prostate; obesity; osteoclast; thymus;
KW   osteoporosis; arthritis; testis; lung; thyroiditis; thyroid; digestion;
KW   endocrine; metabolism; regulation; malabsorption; gastritis; neoplasm.
OS   Homo sapiens.
PN   WO9839446-A2.
PD   11-SEP-1998.
PF   06-MAR-1998; U04492.
PR   06-MAR-1997; US-038621.
PR   07-MAR-1997; US-040161.
PR   07-MAR-1997; US-040162.
PR   07-MAR-1997; US-040163.
PR   07-MAR-1997; US-040333.
PR   07-MAR-1997; US-040334.
PR   07-MAR-1997; US-040336.
PR   07-MAR-1997; US-040626.
PR   11-APR-1997; US-043311.
PR   11-APR-1997; US-043312.
PR   11-APR-1997; US-043313.
PR   11-APR-1997; US-043314.
PR   11-APR-1997; US-043315.
PR   11-APR-1997; US-043568.
PR   11-APR-1997; US-043569.
PR   11-APR-1997; US-043578.
PR   11-APR-1997; US-043579.
PR   11-APR-1997; US-043580.
PR   11-APR-1997; US-043669.
PR   11-APR-1997; US-043670.
PR   11-APR-1997; US-043671.
PR   11-APR-1997; US-043672.
PR   11-APR-1997; US-043674.
PR   23-MAY-1997; US-047492.
PR   23-MAY-1997; US-047500.
PR   23-MAY-1997; US-047501.
PR   23-MAY-1997; US-047502.
PR   23-MAY-1997; US-047503.
PR   23-MAY-1997; US-047581.
PR   23-MAY-1997; US-047582.
PR   23-MAY-1997; US-047583.
PR   23-MAY-1997; US-047584.
PR   23-MAY-1997; US-047585.
PR   23-MAY-1997; US-047586.
PR   23-MAY-1997; US-047587.
PR   23-MAY-1997; US-047588.
PR   23-MAY-1997; US-047589.
PR   23-MAY-1997; US-047590.
PR   23-MAY-1997; US-047592.
PR   23-MAY-1997; US-047593.
PR   23-MAY-1997; US-047594.
PR   23-MAY-1997; US-047595.
PR   23-MAY-1997; US-047596.
PR   23-MAY-1997; US-047597.
PR   23-MAY-1997; US-047598.
PR   23-MAY-1997; US-047599.
PR   23-MAY-1997; US-047600.
PR   23-MAY-1997; US-047601.
PR   23-MAY-1997; US-047612.
PR   23-MAY-1997; US-047613.
PR   23-MAY-1997; US-047614.
PR   23-MAY-1997; US-047615.
PR   23-MAY-1997; US-047617.
PR   23-MAY-1997; US-047618.
PR   23-MAY-1997; US-047632.
PR   06-JUN-1997; US-048964.
PR   06-JUN-1997; US-048974.
PR   06-JUN-1997; US-056631.
PR   06-JUN-1997; US-056632.
PR   22-AUG-1997; US-056636.
PR   22-AUG-1997; US-056637.
PR   22-AUG-1997; US-056662.
PR   22-AUG-1997; US-056664.
PR   22-AUG-1997; US-056845.
PR   22-AUG-1997; US-056862.
PR   22-AUG-1997; US-056864.
PR   22-AUG-1997; US-056872.
PR   22-AUG-1997; US-056874.
PR   22-AUG-1997; US-056875.
PR   22-AUG-1997; US-056876.
PR   22-AUG-1997; US-056877.
PR   22-AUG-1997; US-056878.
PR   22-AUG-1997; US-056879.
PR   22-AUG-1997; US-056880.
PR   22-AUG-1997; US-056881.
PR   22-AUG-1997; US-056882.
PR   22-AUG-1997; US-056884.
PR   22-AUG-1997; US-056886.
PR   22-AUG-1997; US-056887.
PR   22-AUG-1997; US-056888.
PR   22-AUG-1997; US-056889.
PR   22-AUG-1997; US-056892.
PR   22-AUG-1997; US-056893.
PR   22-AUG-1997; US-056894.
PR   22-AUG-1997; US-056903.
PR   22-AUG-1997; US-056908.
PR   22-AUG-1997; US-056909.
PR   22-AUG-1997; US-056910.
PR   22-AUG-1997; US-056911.
PR   05-SEP-1997; US-057650.
PR   05-SEP-1997; US-057650.
PR   05-SEP-1997; US-057761.
PA   (HUMA-) HUMAN GENOME SCI INC.
PI   Bednarik DP, Brewer LA, Carter KC, Duan R, Ebner R, Endress GA,
PI   Feng P, Ferrie AM, Fischer CL, Graves KA, Greene JM, Hu JS,
```

PI Kyaw H, Lafleur DW, Li Y, Moore PA, Ni J, Olsen HS, Rosen CA,
PI Ruben SM, Shi Y, Soppet DR, Young PE, Yu GL, Zeng Z;
DR WPI; 98-609887/51.
DR P-PSDB; W75091.
PT New isolated human genes and the secreted polypeptides they encode
PT - useful for diagnosis and treatment of e.g. cancers, neurological
PT disorders, immune diseases, inflammation or blood disorders
PS Claim 1; Page 190-191; 447pp; English.
CC This sequence represents a nucleic acid molecule which encodes a secreted
CC human protein. The gene number, and the clone it is derived from, are
CC detailed in the descriptor line. The gene can be used to generate fusion
CC proteins by linking to the gene to a human immunoglobulin Fc portion
CC (e.g. V3145) for increasing the stability of the fused protein as
CC compared to the human protein only.
CC The invention relates to 70 novel genes and their fragments (nucleic acid
CC sequences; V3154-V34276; amino acid sequences W75057-W75179) which
CC are useful for preventing, treating or ameliorating medical conditions
CC e.g. by protein or gene therapy. Also, pathological conditions can be
CC diagnosed by determining the amount of the new polypeptides in a sample
CC or by determining the presence of mutations in the new polynucleotides.
CC Specific uses are described for each of the 70 polynucleotides, based on
CC which tissues they are most highly expressed in (see V34154 for described
CC uses).
SQ Sequence 2378 BP; 518 A; 690 C; 662 G; 506 T;

alignment_scores:
         Quality: 221.00   Length: 242
Percent Similarity: Ratio: 1.601   Gaps: 11
                            57.025   Percent Identity: 30.992

Align seg 1/1 to: V34188 from: 1 to: 2378
                   to: V34188

20 ArgArgArgSerGluProSerSerSerSerSerSerSerProAlaAl 36
    ||||||||||||||||::|||
17 CGCCGCGCCGCAGAA.........................TCCACGCTGGCTCCGTGCGC 51

36 aProValaArgProArgProCysProAlaValProAlaProGlyA 53
    |||| |||
52 CATGGTCACCCACAGCAAGTTTCCCGCCCGGGATGAGCCGCCCTGG 101

53 spThrHisPheArg.........ThrPheArgSerHisSerAspTyrArgArg 67
    ||||| |::|||| |||:||||:||||:|:
102 ACACCAGCCTGCCCGCCTCCAAGACCTTCAGCTCCAAGAGGAGTACCAGCGTG 151

68 IleThrArgThrSerAlaLeuLeuAlaCysGlyPheTyrTrpGlyPr 84
    |:::::||||||||||| :::|||::|||||||||||::
152 GTGGTGAACGC.AGTGCGCAAGTGCAGGAGACGGCTTCTACTGGACGGC 200

84 oLeuSerValHisGlyAlaHisGluArgLeuArgAlaGluProValGlyT 101
    ||::: |||::
201 AGTGACCGGCGCGGAGGGAACCTGCTCCTCAGTGCCGAGCCCGCGGCA 250

101 hrPheLeuValArgAspSer.ArgGlnArgAsnCysPhePheAlaLeuSe 117
    |||||||||:::|||||| :::::||||||::|||:||:||
251 CCTTTCTGCAATCCGCGACCAGCTCGGGACGACGCGGCCCACTTCTTCAGCGTCAG 300

117 rValLysMetAlaSerGlyProThrSerIleArgValHisPheGlnAlaG 134
    ||||| :::||:|||||||::|
301 CGTCAAGACCAGTCTGGGACAAGAACCTGCGCATCCAGTGTGAGGGG 350

134 lyArgPheHisLeuAspGlySer........ArgGluThr 144
    ||:|| ||:|||
351 GCAGCTTCTCTCGCAGAGCGATCCCCGGAGCACGCACCCGTGGSCCGGC 400

145 PheAspCysLeuPheGluLeuLeuGluHisTyrVal... 156
    ||||||||:::|||||||||||:::|||||||::::
401 TTCGACTCGTGCTCAACTGGTGCACCACTACATGCCCCGCCCCCTGGAGC 450

156 156

451 CCCCTCCTTCCCCTCGCCACCTACTGAACCGTCCGAGGTGCCCGAGC 500
                                  ::: |||||||||
157 ....AlaAlaProArgArg......... 161

501 AGCCGTCTGCCCAGCCACCTCCCTGGGAGTCCCCAGAAGAGACCTATTAC 550
                              ::: ||||||||
162 ....MetLeuGlyAlaProLeuAr 168

551 ATCTACTCCGGGGGCGGAGGAAGATCCCCCTGGGTGTTGACCGGCCCCTC.. 598
                                     |||||||||||||:::
168 gGlnArgValArgProLeuGlnGluLeuCysGlnArgIleValA 185

599 .TCCTCCAACGTGCGCCACTCTTCAGCAATCTCTGTCGGGAAGACCGTCAACG 647
        ::: ||||||:::||||||:::|||::::
185 laAlaVal...GlyArgGluAsnLeuAlaArgIleProLeuAsnProVal 200

648 GCCACCTGGACTCCTATGAGAAAAGTCACCACCAGCCTGCCG...GGGCCCATT 694
                      |||||||||||||:: :::||||:::
201 LeuArgAspTyrLeuSerSerPhe 208

695 ...CGGGAGGTTCCTGGACCAGTAC 715

seq_documentation_block:
ID V38681 standard; DNA; 848 BP.
AC V38681;
DT 27-OCT-1998 (first entry)
DE Homo sapiens SOCS11 gene.
KW SOCS; suppressor of cytokine signalling; PCR primer;
KW autoimmune disease; diagnosis; cancer; treatment;
KW cytokine mediated cellular responsiveness; hyperimmunity;
KW immunosuppression; allergies; hypertension; ss.
OS Homo sapiens.
FH Key   Location/Qualifiers
FH
FT CDS   1..624
FT   /tag= a
FT   /product= SOCS11 protein
PN WO9820023-A1.
PD 14-MAY-1998.
PF 31-OCT-1997; AU0729.
PR 14-FEB-1997; AU-005117.
PR 01-NOV-1996; AU-003384.
PA (HALL-) HALL INST MEDICAL RES WALTER & ELIZA.
PI Alexander WS, Hilton DJ, Metcalf D, Nicholson SE,
PI Nicola NA, Richardson RT, Starr R, Viney EM, Willson TA;
DR WPI; 98-286654/25.
DR P-PSDB; W62623.
PT Suppressor of cytokine signalling proteins - useful to treat
PT disease; injury or abnormality involving cytokine mediated cellular
PT responsiveness e.g. hyperimmunity, immunosuppression, allergies and
PT hypertension
PS Claim 14; Page 157-158; 325pp; English.
CC The sequence is that of a gene encoding a suppressor of cytokine
CC signalling protein (SOCS). SOCS can be used to screen for naturally
CC occurring antibodies to SOCS, which may occur, e.g. in some autoimmune
CC diseases. Alternatively, specific antibodies can be used to
CC screen for SOCS, which is useful as a knowledge of SOCS levels
CC may be important for the diagnosis of certain cancers. Soluble
CC SOCS polypeptides can be used to treat disease, injury or
CC abnormality involving cytokine mediated cellular responsiveness,
CC e.g. hyperimmunity, immunosuppression, allergies and hypertension.
SQ Sequence 848 BP; 224 A; 215 C; 207 G; 202 T;

alignment_scores:
         Quality: 147.50   Length: 157
Percent Similarity: Ratio: 1.799   Gaps: 2
                            52.229   Percent Identity: 26.752

alignment_block:
US-08-962-560A-4 x V38681

Align seg 1/1 to: V38681   from: 1 to: 848

```
  1 TTGGAAAGTGTGGTTGGTTGATTGGGGGGCCAATGAATTGGGAAGAATGCAGA  50
    |||::|  ||||||||||||||::|||||::|||||||||||||||::  |||:
 75 LeuAspAlaCysGlyPheTyrTrpGlyProLeuSerValHisGlyValAlaHl  91

 51 GATGAAGCTGAAAGGGAAACCAGAAGGGGTTCTTCCCTGGGTACGAGACAGTT 100
    ::||||:  |||||||||::|||||||||||||||::||||||||||||||||
 91 sGlyArgLeuArgAlaGluProValGlyThrPheLeuLeuValArgAspSerA 108

108 rgGlnArgAsnCysPhePheAlaLeuSerValLysMetAlaSerGlyPro 124
    ||||||||||||||||||||||||:||:|||::|||::||||||||||||
108 CTGATCCTCGTTACAATCCTGAGCCCTCAGTTTCCGATCACACAGGTATCACC 150

151 euLeuGluHls........................................ 154
    ::|:|
101 CTGATCCTCGTTACAATCCTGAGCCCTCAGTTTCCGATCACACAGGTATCACC 150
```

seq_documentation_block:

```
ID   V38668 standard; DNA; 1221 BP.
AC   V38668;
DT   27-OCT-1998  (first entry)
DE   Homo sapiens SOCS5 gene.
KW   SOCS; suppressor of cytokine signalling; PCR primer;
KW   autoimmune disease; diagnosis; cancer; treatment;
KW   cytokine mediated cellular responsiveness; hyperimmunity;
KW   immunosuppression; allergies; hypertension; ss.
OS   Homo sapiens
PN   WO982023-A1.
PD   14-MAY-1998.
PR   31-OCT-1997;  AU0729.
PR   14-FEB-1997;  AU-005517.
PR   01-NOV-1996;  AU-003384.
PA   (HALL-) HALL INST MEDICAL RES WALTER & ELIZA.
PI   Alexander WS, Hilton DJ, Metcalf D, Nicholson SE,
PI   Nicola NA, Richardson RT, Starr R, Viney EM, Willson TA;
DR   WPI: 98-286854/25.
PT   Suppressor of cytokine signalling proteins - useful to treat
PT   disease, injury or abnormality involving cytokine mediated cellular
PT   responsiveness e.g. hyperimmunity, immunosuppression, allergies and
PT   hypertension
PS   Disclosure; Page 134-135; 325pp; English.
CC   The sequence is that of a gene encoding a suppressor of cytokine
CC   signalling protein (SOCS). SOCS can be used to screen for naturally
CC   occurring antibodies to SOCS, which may occur, e.g. in some autoimmune
CC   diseases. Alternatively, specific antibodies can be used to
```

```
CC   screen for SOCS, which is useful as a knowledge of SOCS levels
CC   may be important for the diagnosis of certain cancers. Soluble
CC   SOCS polypeptides can be used to treat disease, injury or
CC   abnormality involving cytokine mediated cellular responsiveness,
CC   e.g. hyperimmunity, immunosuppression, allergies and hypertension.
SQ   Sequence 1221 BP; 347 A; 259 C; 275 G; 340 T;
```

alignment_scores:

```
     Quality:       145.00           Length:       142
     Ratio:           1.790          Gaps:           4
     Percent Similarity: 57.042   Percent Identity: 29.577
```

alignment_block:
US-08-962-560A-4 x V38668

Align seg 1/1 to: V38668   from: 1 to: 1221

```
 81 TyrTrpGlyProLeuSerValHisGlyValAlaHisGlyValArgLeuArgAlaGl  97
    |||||||::  |:|| ::  |||  ::|  ||||||::|||::
446 TACTGGGGAGGTGGAACCGGTTATGAAGCAGAAGCCCTTCTCGAAGGGAA 495

 97 uProValGlyThrPheLeuValArgAspSerArgGlnArgAsnCysPhep 114
    ||||  ||||||||||||||::|||||::|||::|||||||||||||::||||
496 ACCTGAAGGCACGTTTTTGCTCAGGGGACTTCGCGCCAAGAGGACTACTTCT 545
```

seq_documentation_block: N_Geneseq_36:V38687

```
ID   V38687 standard; cDNA; 2438 BP.
AC   V38687;
DT   27-OCT-1998  (first entry)
DE   Mus musculus SOCS14 cDNA.
KW   SOCS; suppressor of cytokine signalling; PCR primer;
KW   autoimmune disease; diagnosis; cancer; treatment;
KW   cytokine mediated cellular responsiveness; hyperimmunity;
KW   immunosuppression; allergies; hypertension; ss.
OS   Mus musculus
PN   WO9820023-A1.
PD   14-MAY-1998.
FT   Key             Location/Qualifiers
FT   CDS             2..1630
FT                   /*tag= a
FT                   /product= SOCS14 protein
PN   WO982023-A1.
PD   14-MAY-1998.
PF   31-OCT-1997;  AU0729.
```

PR   14-FEB-1997;
PR   01-NOV-1996; AU-003384.
PA   (HALL-) HALL INST MEDICAL RES WALTER & ELIZA.
PI   Alexander WS, Hilton DJ, Metcalf D, Nicholson SE,
PI   Nicola NA, Richardson RT, Starr R, Viney EM, Willson TA;
DR   WPI: 98-286854/25.
DR   P-PSDB; W62625.
PT   Suppressor of cytokine signalling proteins - useful to treat
PT   disease, injury or abnormality involving cytokine mediated cellular
PT   responsiveness e.g. hyperimmunity, immunosuppression, allergies and
PT   hypertension
PS   Claim 14; Page 165-167; 325pp; English.
CC   The sequence is that of a gene encoding a suppressor of cytokine
CC   signalling protein (SOCS). SOCS can be used to screen for naturally
CC   occurring antibodies to SOCS, which may occur, e.g. in some autoimmune
CC   diseases. Alternatively, specific antibodies can be used to
CC   screen for SOCS, which is useful as a knowledge of SOCS levels
CC   may be important for the diagnosis of certain cancers. Soluble
CC   SOCS polypeptides can be used to treat disease, injury or
CC   abnormality involving cytokine mediated cellular responsiveness,
CC   e.g. hyperimmunity, immunosuppression, allergies and hypertension.
SQ   Sequence 2438 BP; 676 A; 483 C; 556 G; 721 T;

alignment_scores:
         Quality: 138.50   Length: 217
         Ratio: 1.248   Gaps: 9
Percent Similarity: 51.152   Percent Identity: 26.267

alignment_block:
US-08-962-560A-4 x V38687

Align seg 1/1   to: V38687   from: 1 to: 2438

```
     8 AlaAlaAlaAspAsnAlaIleSerProAlaAlaGluProGlnArgArgSerGl  24
       |||   ::||||||||||||||||:|||||:||:::|||||||||||||||||
  1029 GCTCCAGAAAAAGGAATAAGCCCAGGTGGGAAATGGAAGAGGAGATCCTG 1078

    24 uProSerSerSerSerSerSerSerProAlaAlaProVal.ArgPro       40
       ||||||:|||||||||||||||:||||||||||:||||  |||||:|||
  1079 CAGTTGGAGGCACCTCCTAAGTTCCACACCACCAGATCG...ACTACGTCCA 1125

    41 ArgProCysProAlaValProAlaProAlaProGlyAspThrHisPheAr  57
       ||||||||||||      :: |||||||::|||
  1126 CTGCCTTGTTCCANACCTCCT.......... 1146

    57 gThrPheArgSerHisSerAspTyr.ArgArgIleThrArgThrSerAla  73
       |||||||||||||||         ::
  1147 .......TCAGATCAGTAAC..... 1159

    74 LeuLeuAspAlaCysGlyPheThrTrpGlyProLeuSerValHisGlyAl  90
       ||||  ||||||   |||||||||   :::||
  1160 ......AATCCGTGC....TACTGGGGTGTCATGGACAAATATGCAGC 1197

    90 aHisGluArgLeuArgAlaGluProValGlyThrPheLeuValArgAspS 107
       ||||:|||||||||::||||||||||||||||:||::||||||||:|||
  1198 CGAAGCTCTGCTGGAAGGAAAGCCAGGAGGGCACCTTTTTACTTCGAGATT 1247

   107 erArgGlnArgAsnCysPhePheAlaLeuSerValLysMetAlaSerGly 123
       :::  |||||||   |||||||||||   ::::|| ::::||
  1248 CAGCGCAGGAGAATATTAATTCCTCGTTAGTTTAGACCCTACAGTCGT 1297

   124 ProThrSerIleArgValHisPheGlnAlaGlyArgPheHisLeuAspGl 140
       |::||::|||||:||||||||||:|||||||:|||||||||:||
  1298 TCTCTTCATCGCTAGAATTGAGCAGTGGAATCATAACTTAGCTTGATGC 1347

   140 ySerArgGluThrPheAspCysLeuPheGlu.........         151
       ||::||:::|:|||::::                .L.151.
  1348 CCATGAT....CCTTGTGTCTTCCATTCTCCTGATATTACTGGGC 1388

   151 euLeuGluHisTyrValAlaAlaProAlaArgArgMet..........  162
       ||||||||||                                ||| 162
  1389 TCCTGGAACACTATAAGGACCCCAGTGCCTGTATGTTCTTTGAGCCGCTC 1438
```

```
   163 LeuGlyAlaProLeuArgGlnArgArgValArgProLeuGlnGluLeuCy 179
       |||:|||||||||||||||:|||||||||||||:||||:|||::
  1439 TTGTCCACTCCCCTTAATCCGGACGCGTTCCCCTTTCCTTTGGCATATTTG 1488

   179 sArgGlnArgIleValAlaAlaValGlyArgGluAsnLeuArgIleP    196
       |||   ::||| ||||||||||||||||||::||:::||   ::|
  1489 CAGAACGGTTATTTGTTMAATTGTACGACTTACGATGGCATGGATGCCCTTC 1538

   196 roLeuAsnProValLeuArgAspTyrLeuSerSerPheProPheGln    211
       ||:::   |||||||:|||||||||||:|||||||:::::
  1539 CCATTCCTTCGCCTATGAAAATTGTACCTGAAGGAATACCATTATAA 1585
```

seq_name: N_Geneseq.36:V71075

seq_documentation_block:
ID   V71075 standard; cDNA; 2532 BP.
AC   V71075;
DT   08-FEB-1999 (first entry)
DE   Green fluorescent protein-protein tyrosine phosphatase 1C fusion product.
KW   Human; protein tyrosine phosphatase 1C gene; fusion protein;
KW   green fluorescent protein; GFP; intracellular signalling; chimera; ss.
OS   Chimeric - Aequorea victoria.
OS   Chimeric - Homo sapiens.
FH   Key   Location/Qualifiers
FT   CDS   1..2532
FT   /*tag= a
PN   WO9845704-A2.
PD   15-OCT-1998.
PF   07-APR-1998; DK0145.
PR   07-APR-1997; DK-000392.
PA   (NOVO) NOVO-NORDISK AS.
PI   Kasper A, Petersen Bjorn S, Scudder K, Thastrup O,
PI   Tullin S;
DR   WPI: 98-594491/50.
DR   P-PSDB; W85030.
PT   Determining effect on signalling pathways in live cells from
PT   redistribution of luminophores - specifically fusions of green
PT   fluorescent protein with a signalling component, and new apparatus,
PT   particularly for identifying toxins and potential therapeutic agents
PS   Claim 63; Pages 213-217; 326pp; English.
CC   The present sequence encodes a green fluorescent protein (GFP)-human
CC   protein tyrosine phosphatase 1C fusion protein. The fusion protein is
CC   used in an assay to exemplify the invention. The specification describes
CC   how quantitative information about the influence of a molecule on a
CC   cellular response is obtained by recording the variation, caused by
CC   the molecule, on mechanically intact living cells, in the spatially
CC   distributed light emitted from a luminophore present in the cells. The
CC   variation in light emission is processed to provide information that
CC   correlates spatial distribution to the degree of the molecule. The method
CC   is used to identify agents that (in)directly affect intracellular
CC   signalling, especially to screen for potential therapeutic agents or
CC   toxins, and to identify new drug targets.
SQ   Sequence 2532 BP; 613 A; 742 C; 758 G; 419 T;

alignment_scores:
         Quality: 126.50   Length: 147
         Ratio: 1.421   Gaps: 7
Percent Similarity: 60.544   Percent Identity: 34.694

alignment_block:
US-08-962-560A-4 x V71075

Align seg 1/1   to: V71075   from: 1 to: 2532

```
    15 ProAlaAlaGluProArgArgArgSerGluProSerSerSerSerSe  31
       |||||   ::||||||||||  ::||||||:||:::
   549 CCAGAGAACACCCCCATCGGCGA...CGGCCCCGTCGTCCTGCCGGACA 595

    31 rSerSerProAlaAlaProValArgProArgProCysProAlaValProA 48
       |||   ::||||||||:::||||||
   596 ACCACTACCTGAGCACCAGTCCGCCCTGAGCAAAGACCCCAAGAGAAG 645
```

```
 48  laProAlaProGlyAspThrHisPheArg...ThrPheArgSerHisSer  63
     |||  :::  ||||::::||||||||
646  CGCGATCACAATGGGTCCTGGAGTTCGTGACCCGGCCGGGGATCACTCT  695

 64  AspTyrArgArgIle.......ThrArgThrSerAlaLeuLeuAspAla.C  78
     :::  ||:||||||:
696  CGGCATGGACGAGCTGTACAAGTCCGGACTCAGATCTCGAGAGATGCTGT  745

 78  ys...GlyPheTyrTrpGlyProLeuSerValHisGlyAlaHisGluArg  93
     ||:::||  ||||||
746  CCCGTGGGGTGGTTTCACCGAGACCTCAGTGGGGCTGGAATGCAGAGACCCTG  795

 94  LeuArgAlaGluProVal...GlyThrPheLeuValArgAspSerArgGl  109
     |||:::::||||||||||||  ||||::
796  CTCAAGGGCCGAGGTGTCCACGGTAGCTTCCTGGCTCGGCCCAGTCGGCAA  845

109  nArgAsnCysPhePheAlaLeuSerValLysMetAlaSerGlyProThrS  126
     ::|||||||||:||:||:::||||||::
846  GAACCAGGGTGACTTCTCCGCTCTCCGTCAGGGTGGGGATCAGGTGACCC  895

126  erIleArgValHisPheGlnAlaGlyArgPheHisLeuAspGlySerArg  142
     |||||||:::|||  :::  |||:::
896  ATATTCGGATCCAGAACTCAGGGGGATTTCTATGACCTGTATGGAGGG...  942

143  GluThrPheAspCysLeuPheGluLeuLeuGluHisTyr  155
     |||  |||  ||||||:::||:::||:::|||
943  GAGAAGTTTGCGACTCTGACAGAGCCTGGTGGGAGTACTAC  981
```